All India Coordinated Research Project on Agrometeorology (AICRPAM)

ICAR-Central Research Institute for Dryland Agriculture,

Hyderabad, Telangana – 500059

Weather conditions and Agromet advisories issued during May 2021

1. General Weather Conditions

During 01-31 May 2021, the country as a whole received 155.2 mm rainfall, which is 18% excess rainfall, compared to the normal rainfall (131.7 mm) for the same period. Out of 36 meteorological sub-divisions, rainfall was large excess, excess, normal, deficient, large deficient and no rain in 17, 9, 6, 4, 0 and 0 sub-division(s), respectively as depicted in Fig. 1.



@AICRPAM, CRIDA Based on the data given by IMD

2. Agromet advisories issued by AICRPAM centers during May 2021

Date	Advisory issued	Agromet Advisory issued
of issue	for the region	
07-05-2021	Tamil Nadu	Prevailing hot and dry weather condition is favorable for fruit dropping in Acid Lime. Farmers are advised to provide frequent water supply and add a layer mulch under trees.
	Marathwada	Due to the chances of thunderstorm and rainfall, it is advised to harvest mature Mango fruits as early as possible.
	Hisar	Due to the chances of increase in day temperature, farmers are advised to apply light irrigation in planted vegetable crops and fruits plants as per requirement.
	Jammu	Due to chances of rainfall, farmers are advised to postpone the irrigation in vegetable crops.
	Uttarakhand	It is advised to provide mechanical support to the fruit bearing trees and vegetable creepers to prevent from strong winds.
14-05-2021	Madhya Pradesh	Due to hot and dry weather condition, it is advised to sprinkle spray of water on the sheds of poultry and dairy houses to reduce temperature of that area and provide coolers and wet them on the doors of poultry houses to reduce heat inside the house.
	South Interior Karnataka	For the management of incidence of leaf hopper it is advised to spray Azadirachtin (10,000 ppm) @ 7.0 ml/ litre of water.
-	Madhya	To protect fruit crops from high temperature, it is
	Maharashtra	advised to provide shades on fruit crops.
21-05-2021	Kerala	Because of heavy rainfall and flash floods due to 'Tauktae' cyclonic storm, many plantation crops were uprooted. Water stagnation in the pit of coconut seedlings and other plantation crops occurred. Therefore, farmers are advised to prepare proper drainage system and build ridges around the field to drain out excess water from the field. Soil erosion may cause loss of micro nutrients viz. Ca, Mg, B and Zn which can be compensated by application of Sampoorna, Lime, dolomite etc.
	Konkan	Due to cyclonic storm, 'Tauktae', there is breaking, twisting and uprooting of complete plants of mango, cashewnut, spices, coconut and arecanut. Farmers are advised to remove the

1		· · · · · · · · · · · · · · · · · · ·
		dead trees, dried branches and dispose it off, to
		avoid incidence of stem borer in mango and
		cashewnut orchards and red palm weevil in
		coconut and arecanut orchards.
		Because of heavy rainfall with gusty winds due
21-05-2021		to the cyclone 'Tauktae', it is advised to drain
	Gujarat	out excess water from crops field and harvest of
		the mature crops and light sun drying of the
		harvested product to avoid post-harvest loss due
		to infection of fungi/bacteria.
		Due to the possibility of light to moderate
	Chhattisgarh	cloudiness and light showers, farmers are
		advised for harvesting of summer paddy.
		Because of heavy rain, thunderstorm, strong
		wind and high waves due to 'Yaas' cyclonic
	West Bengal	storm, it is advised to harvest the <i>Boro</i> rice, all
	tt est Dengar	the matured pulses and groundnut crops and
		store them at safe places after sun drying.
		Because of heavy rain/thundershower due to the
		cyclone 'Yaas', it is advised to prepare
	Odisha	arrangement for drainage of rainwater from
28-05-2021	Uuisiia	different crop fields like paddy, jute, sugarcane,
		sesamum, vegetables and flowers to avoid
		submergence.
	Vidharkka	Under prevailing weather condition, farmers are
	Vidharbha	advised for deep summer ploughing of harvested
		fields to expose the hibernating insects,
		larvae/pupa to sun and predator birds.
		Due to rainy weather condition, farmers are
		advised to prepare nursery bed for sowing of Sali
	Assam	rice. Recommended varieties are Ranjit, Ranjit
		Sub 1, Bahadur, Satyaranjan, Basundhara,
		Mahsuri, Ketekijoha etc.
	Eastern Uttar	Inter-cultural operation is advised to check the
	Pradesh	evapotranspiration losses and proper utilization
		of nutrient to the crop.